


Accepted Manuscript

Carbonated mantle domains at the base of the Earth's transition zone

Wei-dong Sun, Chris J. Hawkesworth, Chao Yao, Chan-chan Zhang, Rui-fang Huang, Xi Liu, Xin-lei Sun, Trevor Ireland, Mao-shuang Song, Ming-xing Ling, Xing Ding, Zhao-feng Zhang, Wei-ming Fan, Zhong-qing Wu



PII: S0009-2541(17)30430-8
DOI: [doi: 10.1016/j.chemgeo.2017.08.001](https://doi.org/10.1016/j.chemgeo.2017.08.001)
Reference: CHEMGE 18429

To appear in: *Chemical Geology*

Received date: 5 May 2017
Revised date: 31 July 2017
Accepted date: 2 August 2017

Please cite this article as: Wei-dong Sun, Chris J. Hawkesworth, Chao Yao, Chan-chan Zhang, Rui-fang Huang, Xi Liu, Xin-lei Sun, Trevor Ireland, Mao-shuang Song, Ming-xing Ling, Xing Ding, Zhao-feng Zhang, Wei-ming Fan, Zhong-qing Wu, Carbonated mantle domains at the base of the Earth's transition zone, *Chemical Geology* (2017), doi: [10.1016/j.chemgeo.2017.08.001](https://doi.org/10.1016/j.chemgeo.2017.08.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Carbonated mantle domains at the base of the Earth's transition zone

Wei-dong Sun^{1,2,3†}, Chris J. Hawkesworth^{4,5†}, Chao Yao⁶, Chan-chan Zhang^{7,9}, Rui-fang Huang⁹,
Xi Liu⁸, Xin-lei Sun⁹, Trevor Ireland¹⁰, Mao-shuang Song⁹, Ming-xing Ling⁹, Xing Ding⁹, Zhao-
feng Zhang⁹, Wei-ming Fan³, Zhong-qing Wu^{6†}

¹ Center of Deep Sea Research, Institute of Oceanography, Chinese Academy of Sciences, Qingdao 266071, China

² Laboratory for Marine Mineral Resources, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, China

³ CAS Center for Excellence in Tibetan Plateau Earth Sciences, Chinese Academy of Sciences, Beijing, 100101 China;

⁴ Department of Earth & Environmental Sciences, University of St Andrews, North Street, St Andrews KY16 9AL, UK

⁵ School of Earth Sciences, University of Bristol, Wills Memorial Building, Queens Road, Bristol

⁶ School of Earth and Space Sciences, University of Science and Technology of China, Hefei 230026, China;

⁷ University of Chinese Academy of Sciences, Beijing 100049, China;

⁸ Key Laboratory of Orogenic Belts and Crustal Evolution; School of Earth and Space Sciences, Peking University, Beijing 100871, China;

⁹ State Key Laboratory of Isotope Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China.

¹⁰ Research School of Earth Sciences, Australian National University, Canberra 0200, Australia

†Corresponding authors: weidongsun@gig.ac.cn; chris.hawkesworth@st-andrews.ac.uk;
wuzq10@ustc.edu.cn

Abstract: The oxygen fugacity of the upper mantle is 3-4 orders of magnitude higher than that of the lower mantle and this has been attributed to Fe^{2+} disproportionating into Fe^{3+} plus Fe^0 at pressures greater than 24 GPa. The upper mantle might therefore have been expected to have evolved to more oxidizing compositions through geological time, but it appears that the oxygen fugacity of the upper mantle has remained constant for the last 3.5 billion years. Thus, it indicates that the mantle has been actively buffered from the accumulation of Fe^{3+} , and that this is linked to oxidation of diamond to carbonate coupled with reduction of Fe^{3+} to Fe^{2+} . When subducted plates penetrate into the lower mantle, compensational upwelling transports bridgmanite into the transition zone, where it breaks down to ringwoodite and majorite, releasing the ferric iron. The system returns to equilibrium through oxidation of diamond. Early in Earth history, diamond may have been enriched at the base of the transition zone in the Magma Ocean, because it is denser than peridotite melts at depths shallower than 660 km, and it is more buoyant below. Ongoing oxidation of diamond forms carbonate, leading to relatively high carbonate concentrations in the source of ocean island basalts.

Keywords: carbon cycle; carbon concentration; elasticity; first-principles calculations; Earth mantle

1. Introduction

It is generally accepted that the oxygen fugacity of the mantle decreases with increasing depth (Rohrbach and Schmidt 2011), and that mantle plumes originate deep in the Earth's mantle, where carbon is mainly present as diamond (Stagno et al. 2013). Carbonate carried down into the deep mantle through plate subduction also transforms into diamond through a “redox-freezing reaction” (Thomson et al. 2016). The abundance of carbon in the mantle can be estimated from basalts erupting at Earth's surface. Mid-ocean ridge basalt (MORB) is formed in the upper mantle and has carbon concentrations, present as carbonate, of 20 to 300 ppm. The more deeply rooted ocean island basalt (OIB) have higher carbon concentrations (300 to 1300 ppm), suggesting greater carbon contents with increasing depth in the mantle (Dasgupta 2013). The carbon in OIB magmas is present as carbonate, which is seemingly at odds with the decreasing oxygen fugacity with depth. Moreover, the solidus of carbonated peridotite tends to be lower than the current mantle adiabat, except at depths between 300 km to 800 km (Fig. 1a), i.e. carbonate cannot be preserved in the mantle outside this narrow window.

As noted above, diamond is the stable carbon phase in the mantle. If carbon is to change from elemental carbon to carbonate, it must be oxidised (Stagno et al., 2011), and this requires a complementary reduction reaction. The most likely candidate for this reaction is reduction of ferric iron to compensate for the oxidation of carbon. An OIB source with 300 to 1300 ppm C as diamond can be oxidised by ferric Fe of 13-56% for a typical OIB with ca. 8% total iron oxide. This value is about 3-15 times higher than the estimated value ($\text{Fe}^{3+}/\text{Total Fe} < 3\%$) of the upper mantle (Canil and O'Neill 1996). However, the $\text{Fe}^{3+}/\text{Total Fe}$ ratio of OIB is systematically

higher than that in MORB (Evans et al. 2012), even though this appears to contradict the general consensus that the oxygen fugacity of the Earth's mantle decreases with increasing depths.

The lower mantle should be enriched in ferric relative to ferrous iron, because Fe^{2+} disproportionates into Fe^{3+} plus Fe^0 at pressures greater than 24 GPa (Frost *et al.* 2004). Metallic iron will settle down to the core, especially at the time of the Magma Ocean, whereas most of the Fe^{3+} is hosted in bridgmanite with a $\text{Fe}^{3+}/\text{Total Fe}$ ratio of over 0.6 (Frost *et al.* 2004). If Fe^{2+} disproportionation continued through the evolution of the Earth, it would cause oxidation of the whole mantle. Although some authors argued that the oxygen fugacity of the mantle was gradually elevated (Aulbach and Stagno, 2016), Geochemical evidence suggests that the oxygen fugacity of the upper mantle appears to have remained roughly constant over the last 3.5 billion years (Li et al. 2004), suggesting that either such oxidation was restricted to the first billion years (Frost and McCammon 2008), or that the oxygen fugacity in the upper mantle is buffered.

It might be argued that the C contents estimated for the source of OIB reflect a zone of elevated carbon at the base of the transition zone, and that as such they may not be representative of the C content of the lower mantle. If the core contains 1% C, which is an upper estimate of the amount of C in the core (Wood et al. 2013), recent metal-silicate partition coefficients of ~510 to ~5300 for C in the Magma Ocean (Chi et al. 2014) would suggest that there was as little as <2 to <20 parts per million C in the lower mantle. These are up to 3 orders of magnitudes lower than the required C abundance in the sources of OIB. This would suggest that either there is more C in the core and/or lower partition coefficients are applicable. The metal-silicate partition coefficient of C depends strongly on the oxygen fugacity and P-T conditions, and first-principle calculations

show that the metal-silicate partition coefficient of C is 9 ± 3 at 3,200 K and 40 GPa (Zhang and Yin 2012). This would suggest that the C contents of lower mantle rocks in equilibrium with the core are similar to that estimated for the source of OIB.

An alternative possibility is that OIB are derived from a C-rich layer at the base of the transition zone, developed perhaps by diamond segregation at the time of the Magma Ocean. Ongoing oxidation of diamond forms carbonate, leading to the high carbonate concentration in ocean island basalts and explaining the velocity and density anomalies at depths of ~ 660 km.

2. Carbonate layer at the base of the transition zone

We suggest that there is carbonated layer at the base of the transition zone formed through diamond segregation. Diamond is the hardest and thus the least compressible mineral so far recognized. It is also one of the earliest minerals formed in the solar nebula prior to the formation of the Earth and other planets (Hazen et al. 2013), and it is very stable, with high melting temperatures of up to $>5,000$ K (Ogano et al. 2013). Chondritic meteorites have abundant diamond, equivalent to about 750-1500 ppm of carbon (Dai et al. 2002). In contrast to other carbon species such as carbonate and graphite, diamond is stable under the pressures >4 GPa and oxygen fugacity during Fe-Ni precipitation from coexisting silicate melts (Rohrbach and Schmidt 2011; Stagno et al. 2013). Given that the solubility of diamond is very low (~ 2 ppm) in silicate magmas (Hirschmann et al. 2008) and in minerals in the mantle (Wood and Halliday 2005), early diamonds should have been stable in the Magma Ocean and may have pooled together into C-rich domains at the base of the mantle transition zone. Remarkably, diamond is

denser than peridotitic melts in the Earth's mantle at depths above the 660-km discontinuity, but it becomes less dense at greater depths (Figure 1) (Tse and Holzapfel 2008). Consequently, under the conditions of low viscosity and convection in the Magma Ocean, diamond should have accumulated near this density crossover at depths of ~660-km. Such a diamond-rich layer would have been stabilised by the solidification of the Earth, and it would have to have survived the effects of plate subduction and mantle plumes, but it may have been oxidised into a carbonated layer through bridgmanite decomposition.

At least some subducted slabs sink into the lower mantle. In response, lower mantle material, including bridgmanite is forced up into the transition zone through compensational background upwelling. Once in the transition zone, bridgmanite breaks down to ringwoodite and majorite. Ringwoodite accounts for more than 60% of the total silicates, and it has about 5.5% iron. Most of the ferrous iron released from bridgmanite can be hosted in ringwoodite, whereas only a small portion of the ferric iron can be accommodated in majorite. Bridgmanite can have $\text{Fe}^{3+}/\text{Total Fe}$ ratios of 0.6 or higher (Frost *et al.* 2004), whereas majorite has less than ~40% of ferric iron (Fig. 1b) (Rohrbach *et al.* 2007). Therefore, a considerable amount of the ferric iron should be released once bridgmanite breaks down. Consequently, diamond can be oxidized to CO_2 by ferric iron released during the bridgmanite decomposition. Siderite forms through reaction of FeO with CO_2 , and magnesite through reaction of MgO and CO_2 . The material driven out of the lower mantle by the down-going slab drives the mantle in this region to oxidising conditions, and this can be maintained in a quasi-equilibrium state, at least at depths of 300 to 800 km. These carbonate-enriched mantle domains may have acted as the main carbonate supplier to mantle

plumes and probably also to kimberlites. This is one way to explain the systematically higher carbon contents and oxygen fugacity of OIB compared to MORB.

3. Calculation details and results

3.1 The density of diamond

The density profiles of diamond along the geotherm of the normal mantle (Jackson 1998) were calculated by using the third-order high- T Birch-Murnaghan equation of state, following:

$$P = \frac{3}{2} K_{T,0} \left[\left(\frac{\rho(T,P)}{\rho(T,0)} \right)^{\frac{7}{3}} - \left(\frac{\rho(T,P)}{\rho(T,0)} \right)^{\frac{5}{3}} \right] \times \left\{ 1 - \frac{3}{4} (4 - K'_{T,0}) \left[\left(\frac{\rho(T,P)}{\rho(T,0)} \right)^{\frac{2}{3}} - 1 \right] \right\} \quad (1),$$

$$K_{T,0} = K_{300,0} + \left(\frac{\partial K_{T,0}}{\partial T} \right)_P (T - 300) \quad (2),$$

$$\rho(T,0) = \rho_0 / \exp \int_{300}^T \alpha_{T,0} dT \quad (3),$$

$$K'_{T,0} = K'_{300,0} \quad (4),$$

and

$$\alpha_{T,0} = a + bT + c/T^2 \quad (5),$$

where $K_{T,0}$, $K'_{T,0}$, $\rho(T,0)$, and $\alpha_{T,0}$ are the isothermal bulk modulus, its pressure derivative, density, and thermal expansivity of diamond at T and ambient P , respectively. The data used in the calculation were from Reeber and Wang (1996) (Table 1)

3.2 Density and elasticity of magnesite

First principle calculations were performed using the open-source Quantum ESPRESSO package (Giannozzi et al. 2009) based on the density functional theory (DFT) with the local density approximation and a 70 Ry cut-off energy for the plane wave basis. The details on the pseudopotentials for Mg, O, and C can be found in Wang et al. (2017). The structure at certain pressures was optimized using the variable cell-shape damped molecular dynamic (Wentzcovitch et al. 1993). The dynamical matrices on a 4×4×4 mesh for the optimized structure, computed using density functional perturbation theory (DFPT) (Baroni et al. 2001), were interpolated in a regular 12·12·12 mesh to obtain the vibrational density of state.

The Helmholtz free energy in the quasi-harmonic approximation (QHA) is given by

$$F(V,T) = U_o(V) + \frac{1}{2} \sum_{q,m} \hbar \omega_{q,m}(V) + k_B T \sum_{q,m} \ln \left\{ 1 - \exp \left[-\frac{\hbar \omega_{q,m}(V)}{k_B T} \right] \right\} \quad (6),$$

where ω is vibrational frequency, q is the phonon wave vector, m is the normal mode index. The calculated Helmholtz free energies versus volume were fitted by isothermal third-order finite strain equations of state (Eq. 1). The resulting pressure-volume relations are described by a third-order Birch-Murnaghan equation of states (EOS).

The static elastic constant tensors were calculated from stresses generated by small deformation (1%) of the optimized unit cell with internal atomic coordinated relaxation. The Wu and Wentzcovitch (2011) method, which reduces the computation time to a tenth of conventional methods, was used to obtain the high-temperature and -pressure elastic tensors. As shown in Figs. 2-3, the calculated elasticity, density and sound velocities of magnesite agree well with

available experimental data (Chen et al. 2006; Fiquet and Reynard 1999; Hearmon et al. 1979; Hubert et al. 1972; Sanchez-Valle et al. 2001; Yang et al. 2014).

3.3 Density and elasticity of siderite

Previous studies have shown that the density and elasticity of forsterite, its polymorphs, and MgO obtained using the method also agree well with experimental data (Wu and Wentzcovitch 2011; Nunez-Valdez et al. 2012; 2013). Using these calculated data, we found that these mantle minerals show similar temperature and pressure dependence on density and sound velocities. The density to velocity ratios among these minerals vary by less than 5% in most cases of temperature and pressure range of upper mantle (Fig. 4). By assuming this relation for magnesite and siderite, we obtained the density and velocity profiles of siderite along the geotherm (Table 2).

4. Discussion

4.1 Disturbance of the diamond-rich layer

Both crystallization and cooling have influences on the density crossover. Given that the composition of the mantle peridotite is close to that of the peridotite melt, the composition is not supposed to change much during cooling in the early history of the Earth. Once the Earth is solidified, the density crossover change to 410 km (Figure 5). Nevertheless, diamond cannot

move once the viscosity increases to a certain point. Therefore, the diamond layer cannot change dramatically.

Diamond segregation/accumulation driven by differences in density should have stopped after the solidification of the Magma Ocean. Nevertheless, Proterozoic eclogitic diamond (Shirey et al. 2002) indicates that diamond continued to crystallise at least in part in response to the recycling of carbon through subduction zones. This is supported by isotope ratios of diamond and ultrahigh pressure inclusions within it (Walter et al. 2011).

A diamond enriched layer that formed during the Magma Ocean period will subsequently have been disturbed by mantle plumes and plate subduction. Plume magmas that erupted over Earth history could have covered the Earth's surface ~2-4 times over, assuming an eruption rate of 2 km³ per year, and an average thickness of 5-10 km for the volcanic rocks. The diameter of a mantle plume may increase by over 10 times from the base of the transition zone to the Earth's surface. Therefore, less than 5% of a diamond layer at the base of the transition zone might be expected to have been disturbed by mantle plumes over the history of the Earth. Considering the more intense magmatism in the early history of the Earth, this number may perhaps be doubled. Similarly, subducting plates may have disturbed 20-40% of the diamond layer, assuming that plate subduction started at the beginning of the Earth, that all subducted plates penetrate into the lower mantle, they are 50 thousand kilometers long and 90 kilometers thick (crust + oceanic lithosphere mantle), and that on average each plate survives for 100-200 million years. Such necessarily simplistic calculations suggest that there might be at least 50% of a diamond-enriched layer still present at the base of the transition zone.

4.2 Bridgmanite decomposition and oxidation of diamond

Given that at pressures higher than 7 GPa, pyroxene progressively dissolves in garnet (Equation 7), ringwoodite and majorite are the only two silicate minerals at the base of the transition zone. During plate tectonics, subducting oceanic slabs may penetrate into the lower mantle. In response, bridgmanite rises to the transition zone through compensational background upwelling, and breaks down to ringwoodite and majorite. The reaction can be described as Equation (8): ferric iron goes only into majorite, whereas ferrous iron is hosted by ringwoodite, and majorite solid solution (Rohrbach et al. 2007).



Therefore, a considerable proportion of the ferric iron should be released from silicates (Equation 2), and even assuming that all the ferrous iron goes into majorite, one third of the ferric iron is released. Given that the oxygen fugacity of the upper mantle remained constant for the last 3.5 billion years, the implication is that there is an oxygen fugacity buffer at the base of the transition zone. The diamond-carbonate buffer is the most likely candidate.

Diamond at the base of the transition zone is the first to be oxidized to CO₂ by ferric iron released during bridgmanite decomposition (Equation 9), forming siderite and magnesite. These carbonate-enriched mantle domains may then have acted as the main carbonate supplier to mantle plumes and probably also to kimberlites.

4.3 Carbonate and velocity and density anomalies

The density and velocity of the Earth's mantle increase at slower rates at depths between 600-660 km than at shallower depths (Fig. 5) (Dziewonski and Anderson 1981). This is not consistent with nearly linear increases in the velocities and densities of ringwoodite and majorite with increasing depth. In principle, such anomalies either suggest that there are additional phase transitions that have not yet been identified, or there are compositional anomalies. Considering the number of high-pressure experiments that have been performed, a major phase transition of mantle silicate minerals at depths of 600-660 km is unlikely. Instead the geophysical observations are taken to favor a more or less continuous carbonated layer at the base of the Earth's transition zone.

Carbonated domains also plausibly explain the density and velocity profiles between 600-660 km (Fig. 5) (Dziewonski and Anderson 1981). Our first-principle calculations show that magnesite has density and velocity considerably lower than silicates (Fig. 5 and Table 2). Siderite is denser than magnesite, while its velocity is slower than that of magnesite (Sanchez-Valle et al. 2011). The density and velocity profiles are consistent with the addition of up to ~1.3% and ~0.44% of

carbon in the forms of magnesite and siderite, at 660 km, respectively (Table 3). This corresponds to a 60 km thick carbonate-bearing layer with an average carbon content of ~0.87%.

Background upwelling of bridgmanite is taken to be a long term process linked to plate subduction. Therefore, carbonate has likely formed in response to the decomposition of bridgmanite since the onset of plate tectonics. The solidus line of carbonated peridotite is well above the current mantle adiabat, but it is lower than the Archean mantle adiabat, at depths of 600-660 km (Fig. 1a) (Dasgupta 2013), suggesting that such carbonated peridotite may only have been stable in the mantle once the mantle is cold enough, i.e., in the post-Archean era.

A recent study indicates that the transition zone may at least locally have water contents of ~1 wt%, which is much higher than previously recognized (Pearson et al., 2014). The presence of water in ringwoodite would dramatically reduce its density and velocity (Mao et al. 2012), and consequently less carbonate would be required to explain the anomalies at the base of the transition zone. However, the water content in the transition zone is still highly debated, and an additional mechanism would be needed to explain why water is restricted at depths of 600-660 km.

In contrast to carbonates, the velocity of diamond is far higher than those of ringwoodite and majorite. The negative velocity anomaly at the bottom of the Earth's transition zone (Dziewonski and Anderson 1981) therefore suggests that only a small amount of diamond remains in the carbonated layer.

4.4 Carbonate and concentrations of atmospheric CO₂

Carbonate may be subducted into the mantle (Stagno et al., 2015), such that is an integral part of carbon recycling, and there appears to be two pathways for carbon to be recycled into the mantle. There is a shallow level cycle in which carbonate is subducted and remobilized through destructive plate margin magmatism (Li et al. 2017; Zhang et al., 2017; Sun et al., 2017). In a deeper level cycle, subducted carbon transforms into diamond on its way down to the lower mantle, and it may stay in the ultra-low velocity zone or very low velocity provinces above the core-mantle boundary for billions of years together with subducted oceanic crust.

Experiments however show that carbonate-induced melting can occur in deeply subducted lithosphere at near-adiabatic temperatures in the Earth's transition zone and lower mantle (Rohrbach and Schmidt 2011). Mineral segregation in these low velocity regions is likely to occur when there are relatively high degrees of partial melting (Sun et al. 2011). Diamond is much less dense than the surrounding mantle and silicate melts at depths below the 660-km discontinuity, such that diamond in subducted oceanic slabs may be released and carried upwards by mantle plumes.

Ferromagnesian carbonates in the carbonated layer may also be transported up in upwelling mantle domains, e.g., plumes, kimberlites. A high carbonate concentration in the carbonated layer would dramatically lower the mantle solidus, and promote partial melting within mantle plumes. The degree of partial melting of the carbonated layer critically depends on the temperature of the mantle plumes and mantle domains it has sampled. Large igneous provinces

associated with plume heads may be expected to carry up and release more carbon dioxide to Earth's atmosphere. This offers one explanation for the dramatic fluctuation of concentrations of atmospheric carbon dioxide and mass extinctions associated with mantle plumes (Courtillot and Olson 2007; Sobolev et al. 2011).

5. Concluding remarks

Ocean island basalts have systematically higher and more varied carbon contents than mid-ocean ridge basalts. At the depths from which the sources of OIB were derived, the mantle is reducing and most of the carbon is therefore likely to be present in diamond. Yet the oxygen fugacity of OIB is systematically higher than that of mid-ocean ridge basalts, which requires additional oxygen. One way is through the presence of carbonates rather than diamond in the source of OIB. We argue that OIB sampled carbonated domains at the base of the transition zone formed through oxidation of diamond by ferric iron released by breakdown of bridgmanite. Bridgmanite is the main mineral of the lower mantle, which contains up to 60% of ferric iron. When subducted plates penetrate into the lower mantle, compensational background upwelling takes bridgmanite into the transition zone, which transforms into ringwoodite and majorite, releasing ferric iron. This oxidizes diamond, forming carbonated domains along the base of the transition zone, which is then taken to be responsible for both the high carbon contents and oxygen fugacities of OIB. Diamond may be enriched at the base of the transition zone, because it is denser than peridotitic melt at depths of <660 km and becomes less dense at deeper depths, such that it should have accumulated near this density crossover in the Magma Ocean period. The density and seismic velocity anomalies at depths between 600-660 km can also be plausibly

explained by the presence of carbonated peridotitic rocks, suggesting that carbonated domains might form a more or less continuous layer. More high pressure experiments on the densities and velocities of carbonates and carbonated mantle peridotite under the P-T conditions at the base of the transition zone are required for future studies.

Acknowledgements

This study was supported by National Key R&D Program of China 2016YFC0600408, CAS XDB18020000, NSFC 91328204 to W.D.S., State Key Development Program of Basic Research of China (2014CB845905), NSFC 41274087 to Z.Q.W. 41090371 to X.L. Helpful discussions with Charles Langmuir and Mike Walter are highly appreciated. TRI acknowledges a Chinese Academy of Sciences President's International Fellowship for visiting scientists Grant No. 2015VEA003.

References:

- Aulbach, S., Stagno, V., 2016. Evidence for a reducing Archean ambient mantle and its effects on the carbon cycle. *Geology*, 44(9): 751-754.
- Canil, D. & O'Neill, H. S. C. (1996) Distribution of ferric iron in some upper-mantle assemblages. *J Petrol* **37**, 609-635.
- Chen, P.-F., Chiao, L.-Y., Huang, P.-h., Yang, Y.-j., Liu, L.-g., 2006. Elasticity of magnesite and dolomite from a genetic algorithm for inverting Brillouin spectroscopy measurements. *Physics of the Earth and Planetary Interiors* 155, 73-86.

- Chi, H., Dasgupta, R., Duncan, M. S. & Shimizu, N. (2014) Partitioning of carbon between Fe-rich alloy melt and silicate melt in a magma ocean - Implications for the abundance and origin of volatiles in Earth, Mars, and the Moon. *Geochim Cosmochim Acta* **139**, 447-471, doi:DOI 10.1016/j.gca.2014.04.046.
- Courtillot, V. & Olson, P. (2007) Mantle plumes link magnetic superchrons to phanerozoic mass depletion events. *Earth Planet Sc Lett* **260**, 495-504, doi:DOI 10.1016/j.epsl.2007.06.003.
- Dai, Z. R. *et al.* (2002) Possible in situ formation of meteoritic nanodiamonds in the early Solar System. *Nature* **418**, 157-159, doi:Doi 10.1038/Nature00897.
- Dasgupta, R. (2013) Ingassing, storage, and outgassing of terrestrial carbon through geologic time. *Reviews in Mineralogy & Geochemistry* **75**, 183-229.
- Dasgupta, R. & Hirschmann, M. M. (2010) The deep carbon cycle and melting in Earth's interior. *Earth Planet Sc Lett* **298**, 1-13, doi:10.1016/j.epsl.2010.06.039.
- Dziewonski, A. M. & Anderson, D. L. (1981) Preliminary reference Earth model. *Phys Earth Planet In* **25**, 297-356.
- Evans, K. A., Elburg, M. A. & Kamenetsky, V. S. (2012) Oxidation state of subarc mantle. *Geology* **40**, 783-786, doi:Doi 10.1130/G33037.1.
- Fiquet, G., Reynard, B., 1999. High-pressure equation of state of magnesite: New data and a reappraisal. *American Mineralogist* **84**, 856-860.
- Frost, D. J. & McCammon, C. A. (2008) The redox state of Earth's mantle. *Annu Rev Earth Pl Sc* **36**, 389-420, doi:DOI 10.1146/annurev.earth.36.031207.124322.
- Frost, D. J. *et al.* (2004) Experimental evidence for the existence of iron-rich metal in the Earth's lower mantle. *Nature* **428**, 409-412, doi:Doi 10.1038/Nature02413.

- Giannozzi, P. *et al.* (2009).QUANTUM ESPRESSO: a modular and open-source software project for quantum simulations of materials. *Journal of Physics-Condensed Matter* **21**, doi:Artn 395502. Doi 10.1088/0953-8984/21/39/395502
- Hazen, R. M., Downs, R. T., Kah, L. & Sverjensky, D. (2013) Carbon mineral evolution. *Reviews in Mineralogy & Geochemistry* **75**, 79-107.
- Hearmon, R., 1979. The third-and higher-order elastic constants. Landolt-Börnstein Zahlenwerte und Funktionen aus Naturwissenschaften und Technik (Numerical Values and Functional Relationships in Science and Technology), New Series, Group III, 245-286.
- Hirschmann, M. M. & Withers, A. C. (2008) Ventilation of CO₂ from a reduced mantle and consequences for the early Martian greenhouse. *Earth Planet Sc Lett* **270**, 147-155, doi:10.1016/j.epsl.2008.03.034.
- Hoffman, P. F., Kaufman, A. J., Halverson, G. P. & Schrag, D. P. A (1998) Neoproterozoic snowball earth. *Science* **281**, 1342-1346, doi:DOI 10.1126/science.281.5381.1342.
- Humbert, P., Plicque, F., 1972. Elastic Properties of Monocrystalline Rhombohedral Carbonates - Calcite, Magnesite, Dolomite. *Comptes Rendus Hebdomadaires Des Seances De L Academie Des Sciences Serie B* 275, 391-
- Jackson, I. Elasticity, composition and temperature of the Earth's lower mantle: a reappraisal. *Geophys J Int* **134**, 291-311, doi:DOI 10.1046/j.1365-246x.1998.00560.x (1998).
- Li, S.G. et al., 2017. Deep carbon cycles constrained by a large-scale mantle Mg isotope anomaly in eastern China. *National Science Review*, 4(1): 111-120.
- Li, Z. X. A. & Lee, C. T. A. (2004) The constancy of upper mantle fO(2) through time inferred from V/Sc ratios in basalts. *Earth Planet Sc Lett* **228**, 483-493, doi:10.1016/j.epsl.2004.10.006.

- Mao, Z. *et al.* (2012) Sound velocities of hydrous ringwoodite to 16 GPa and 673 K. *Earth Planet Sc Lett* **331**, 112-119, doi:DOI 10.1016/j.epsl.2012.03.001.
- Núñez Valdez, M., Wu, Z., Yu, Y.G., Revenaugh, J., Wentzcovitch, R.M., (2012) Thermoelastic properties of ringwoodite ($\text{Fe}_x\text{Mg}_{1-x}\text{SiO}_4$): Its relationship to the 520km seismic discontinuity. *Earth and Planetary Science Letters* 351-352, 115-122.
- Núñez Valdez, M., Wu, Z., Yu, Y. G. & Wentzcovitch, R. M. (2013) Thermal elasticity of ($\text{Fe}_x\text{Mg}_{1-x}\text{SiO}_4$) olivine and wadsleyite. *Geophys Res Lett* **40**, 290-294, doi:Doi 10.1002/Grl.50131.
- Oganov, A. R., Hemley, R. J., Hazen, R. M. & Jones, A. P. (2013) Structure, bonding, and mineralogy of carbon at extreme conditions. *Reviews in Mineralogy & Geochemistry* **75**, 47-77.
- Ohtani, E. (2009) Melting relations and the equation of state of magmas at high pressure: Application to geodynamics. *Chem Geol* **265**, 279-288, doi:DOI 10.1016/j.chemgeo.2009.04.004.
- Pearson, D. G. *et al.* (2014) Hydrous mantle transition zone indicated by ringwoodite included within diamond. *Nature* **507**, 221-+, doi:Doi 10.1038/Nature13080.
- Reeber, R. R. & Wang, K. Thermal expansion, molar volume and specific heat of diamond from 0 to 3000K. *J Electron Mater* **25**, 63-67, doi:Doi 10.1007/Bf02666175 (1996).
- Rohrbach, A. & Schmidt, M. W. (2011) Redox freezing and melting in the Earth's deep mantle resulting from carbon-iron redox coupling. *Nature* **472**, 209-212, doi:10.1038/nature09899.
- Rohrbach, A. *et al.* (2007) Metal saturation in the upper mantle. *Nature* **449**, 456-458, doi:10.1038/nature06183.

- Sanchez-Valle, C., Ghosh, S. & Rosa, A. D. (2011) Sound velocities of ferromagnesian carbonates and the seismic detection of carbonates in eclogites and the mantle. *Geophys Res Lett* **38**, doi:10.1029/2011gl049981.
- Shirey, S. B. *et al.* (2002) Diamond genesis, seismic structure, and evolution of the Kaapvaal-Zimbabwe craton. *Science* **297**, 1683-1686, doi:DOI 10.1126/science.1072384.
- Sobolev, S. V. *et al.* (2011) Linking mantle plumes, large igneous provinces and environmental catastrophes. *Nature* **477**, 312-U380, doi:Doi 10.1038/Nature10385.
- Stagno, V., Ojwang, D. O., McCammon, C. A. & Frost, D. J. (2013) The oxidation state of the mantle and the extraction of carbon from Earth's interior. *Nature* **493**, 84-+, doi:Doi 10.1038/Nature11679.
- Stagno, V., Frost, D.J., McCammon, C.A., Mohseni, H., Fei, Y., 2015. The oxygen fugacity at which graphite or diamond forms from carbonate-bearing melts in eclogitic rocks. *Contributions to Mineralogy and Petrology*, 169(2).
- Stagno, V. *et al.*, 2011. The stability of magnesite in the transition zone and the lower mantle as function of oxygen fugacity. *Geophysical Research Letters*, 38.
- Sun, W.D. *et al.*, 2011. The fate of subducted oceanic crust: a mineral segregation model. *International Geology Review*, 53(8): 879-893.
- Sun, W.-d., Zhang, C.-c., 2017. Carbon recycling through alkali basalts. *Solid Earth Sciences*, 2(2): 41-42.
- Thomson, A. R., Walter, M. J., Kohn, S. C. & Brooker, R. A. (2016) Slab melting as a barrier to deep carbon subduction. *Nature* **529**, 76-79, doi:doi:10.1038/nature16174.
- Tse, J. S. & Holzapfel, W. B. (2008) Equation of state for diamond in wide ranges of pressure and temperature. *J. Appl. Phys.* **104**, doi:10.1063/1.2969909.

- Wang, W., Qin T., Zhou, C., Wu, Z., Huang, F., (2017) Concentration effect on equilibrium fractionation of Mg-Ca-C-O isotopes in carbonate minerals: insights from first-principles calculations. *Geochim Cosmochim Ac.* 208, 185–197
- Wentzcovitch, R. M., Martins, J. L. & Price, G. D. (1993) Abinitio Molecular-Dynamics with Variable Cell-Shape - Application to MgSiO_3 . *Phys Rev Lett* **70**, 3947-3950, doi:DOI 10.1103/PhysRevLett.70.3947.
- Wood, B. J. & Halliday, A. N. (2005) Cooling of the Earth and core formation after the giant impact. *Nature* **437**, 1345-1348, doi:10.1038/nature04129.
- Wood, B. J., Li, J. & Shahr, A. (2013) Carbon in the core: Its influence on the properties of core and mantle. *Reviews in Mineralogy & Geochemistry* **75**, 231-250.
- Wu, Z. Q. & Wentzcovitch, R. M. (2011) Quasiharmonic thermal elasticity of crystals: An analytical approach. *Phys Rev B* **83**, doi:Artn 184115, Doi 10.1103/Physrevb.83.184115.
- Yang, J., Mao, Z., Lin, J.-F. & Prakapenka, V. B. (2014) Single-crystal elasticity of the deep-mantle magnesite at high pressure and temperature. *Earth Planet Sc Lett* **392**, 292-299, doi:http://dx.doi.org/10.1016/j.epsl.2014.01.027.
- Zhang, Y. G. & Yin, Q. Z. (2012) Carbon and other light element contents in the Earth's core based on first-principles molecular dynamics. *P Natl Acad Sci USA* **109**, 19579-19583, doi:DOI 10.1073/pnas.1203826109.
- Walter, M. J. *et al.* (2011) Deep Mantle Cycling of Oceanic Crust: Evidence from Diamonds and Their Mineral Inclusions. *Science* **333**, 54-57, doi:DOI 10.1126/science.1209300.
- Zhang, G.L., Chen, L.H., Jackson, M.G., Hofmann, A.W., 2017. Evolution of carbonated melt to alkali basalt in the South China Sea. *Nature Geoscience*, 10(3): 229-U105.

Figure Captions

Figure 1. (A) Solidus of carbonated peridotite (Dasgupta and Hirschmann 2010). Carbonated peridotite is stable between ~300-800 km with a post-Archean geotherm (Stagno et al. 2013). During plate subduction, bridgmanite breaks down to ringwoodite and majorite at the base of the transition zone due to compensational background upwelling, releasing ferric iron that oxidizes diamond, forming carbonated mantle domains. These carbonated domains may be sampled by mantle plumes, but not MORB, such that OIB has systematically higher CO₂ than MORB. (B) Proportions of ferric iron in majorite (Rohrbach et al. 2007) and bridgmanite (Frost et al. 2004) under different pressures. Bridgmanite has systematically high proportions of ferric iron. When bridgmanite is broken down during background upwelling across the upper-lower mantle boundary, a portion of the ferric iron is absorbed by majorite, the rest is released, oxidizing diamond to carbon dioxide.

Figure 2. Pressure dependence of the elastic constants of magnesite at 300K. The experimental data are shown in scattered points

Figure 3. Pressure dependence of the sound velocities and density of magnesite at 300K. The experimental data are shown in scattered points

Figure 4. (A) P-wave velocity ratio, (B) density ratio, and (C) S wave velocity ratio among MgO, forsterite, and its polymorphs along the geotherm. The ratios were normalized at ambient conditions. The elastic data for forsterite, wadsleyite, ringwoodite, and mgo are from Wu et al., (2011), Núñez Valdez et al., (2012, 2013).

Figure 5. (A) and (B) Density profiles of magnesite, diamond (Tse and Holzapfel 2008), Preliminary Reference Earth Model (PREM) (Dziewonski and Anderson 1981), peridotitic melt (Ohtani 2009). Diamond in the Magma Ocean is predominately concentrated at the density

crossover between diamond and peridotitic melt near the 660-km discontinuity, and then oxidized to ferromagnesian carbonates by Fe^{3+} released due to background upwelling of bridgmanite. At depths of 660 km, the density of PREM is $3.99 \cdot 10^3 \text{ kgm}^{-3}$, whereas extrapolation of the density curve from 600 km to 660 km gives a density of $4.05 \cdot 10^3 \text{ kgm}^{-3}$. The density anomaly at 660 km depth can be plausibly explained by addition of ~1.1% carbon in the form of magnesite and siderite (Table 3). **(C) and (D)** Velocity profiles of Preliminary Reference Earth Model (PREM), showing anomalies at depths between 600-660 km. Also shown is the velocity of magnesite (this study) based on first-principle calculations. The velocity anomalies of PREM can also be explained by carbonated peridotite (Table 3).




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

Table 1 Density profile of diamond along the normal mantle geotherm.

Pressure (GPa)	Temperature (K)	V/V ₀	density (g/cm ³)
0	1605	1.02	3.53
1	1616	1.01	3.48
2	1627	1.01	3.49
3	1637	1.01	3.50
4	1648	1.01	3.50
5	1658	1.01	3.51
6	1668	1.00	3.52
7	1678	1.00	3.53
8	1688	1.00	3.53
9	1698	1.00	3.54
10	1708	1.00	3.55
15	1755	0.99	3.58
20	1798	0.98	3.61
25	1839	0.97	3.65
30	1877	0.96	3.68
35	1911	0.95	3.71

Table 2 Density and velocities of magnesite and siderite at 660 km and 1826 K.

	Magnesite	Siderite		
		f=1.0	f=0.9	f=1.1
Density (g/cm ³)	3.401	4.401	4.890	4.001
V _S (km/s)	5.234	3.732	4.147	3.393
V _P (km/s)	9.742	7.864	8.738	7.149

$f = \frac{A^{mag}(660km, 1826K)}{A^{sid}(660km, 1826K)} \cdot \frac{A^{sid}(0km, 300K)}{A^{mag}(0km, 300K)}$, where A is density or velocities of magnesite (mag) and siderite (sid). Siderite data at ambient condition is from Sanchez-Valle, et al (2011)

Table 3 The density and velocities for the mantle with best fitting concentration of magnesite and siderite at 660 km and 1826 K.

	Real mantle	f=1	f=0.9	f=1.1
C in magnesite		1.7%	2.3%	1.3%
C in siderite		0.46%	0.50%	0.44%
Density (g/cm ³)	3.99	3.989	3.987	3.990
V _S (km/s)	5.56	5.548	5.544	5.552
V _P (km/s)	10.25	10.259	10.263	10.256

Highlights

Diamond accumulated at ~ 660-km depth in Magma Ocean because of density crossover.

Diamond is oxidised by Fe^{3+} from bridgmanite decomposition, forming carbonated domain.

The carbonated domain can explain higher oxygen fugacity of OIB than that of MORB

The carbonated domain can explain high carbonate concentrations of OIB

The carbonated domain can explain density and velocity anomalies at depths 600-660 km